



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The city as a refuge for insect pollinators

Citation for published version:

Stone, G, Hall, DM, Camilo, GR, Tonietto, RK & Ahrne, K 2016, 'The city as a refuge for insect pollinators', *Conservation biology*. <https://doi.org/10.1111/cobi.12840>

Digital Object Identifier (DOI):

[10.1111/cobi.12840](https://doi.org/10.1111/cobi.12840)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Conservation biology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



The city as a refuge for insect pollinators

Damon M. Hall,
Saint Louis University, Center for Sustainability

Gerardo R. Camilo,
Saint Louis University, Department of Biology

Rebecca K. Tonietto,
David H. Smith Conservation Research Program, Society for Conservation Biology; Saint Louis
University, Department of Biology

Karin Ahrné,
Swedish University of Agricultural Sciences, Swedish Species Information Centre-ArtDatabanken

Mike Arduser,
Missouri Department of Conservation (retired)

John S. Ascher
National University of Singapore, Department of Biological Science

Katherine C. R. Baldock,
University of Bristol, Biological Sciences & Cabot Institute

Robert Fowler,
University of Sussex, School of Life Sciences

Gordon Frankie,
University of California Berkeley, Department of Environmental Science, Policy, & Management

Dave Goulson,
University of Sussex, School of Life Sciences

Bengt Gunnarsson,
University of Gothenburg, Department of Biological and Environmental Sciences

Mick E. Hanley,
Plymouth University, School of Biological Sciences

Janet I. Jackson,
University of Northampton, Department of Environmental and Geographical Sciences

Gail Langellotto,
Oregon State University, Department of Horticulture

David Lowenstein,
University of Illinois-Chicago, Department of Biological Sciences

Emily S. Minor,
University of Illinois-Chicago, Department of Biological Sciences

Jeff Ollerton,
University of Northampton, Department of Environmental and Geographical Sciences

Stacy M. Philpott,
University of California, Santa Cruz, Environmental Studies

Simon G. Potts,
Reading University, Centre for Agri-Environmental Research, School of Agriculture, Policy and
Development

Muzafar H. Sirohi,
University of Northampton, Department of Environmental and Geographical Sciences

Edward M. Spevak,
Saint Louis Zoo, WildCare Institute Center for Native Pollinator Conservation, IUCN SSC Bumblebee
Specialist Group

Graham N. Stone,
The University of Edinburgh, Institute of Evolutionary Biology

Caragh G. Threlfall,
The University of Melbourne, Green Infrastructure Research Group

Abstract:

Urban ecology research is changing how we view the biological value and ecological importance of cities. Lagging behind this revised image of the city are natural resource management agencies' urban conservation programs that historically have invested in education and outreach rather than programs designed to achieve high-priority species conservation results. This essay synthesizes research on urban bee species diversity and abundance to suggest how urban conservation can be repositioned to better align with a newly unfolding image of urban landscapes. We argue that pollinators put high-priority and high-impact urban conservation within reach. In a rapidly urbanizing world, transforming how environmental managers view the city can improve citizen engagement while exploring more sustainable practices of urbanization.

Keywords:

Urban ecology, Pollinator, Insects < Animals, Communications, Conservation planning, Governance, Funding and philanthropy, Politics and policy

Natural resource management¹ (NRM) investments in urban conservation programming are largely aimed at connecting people to nature, rather than efforts believed to achieve high conservation impact. Historically, urban conservation directives have sought to garner broad public support by funding outreach, recreation facilities, and education rather than high-priority conservation outcomes (USFWS, 2015; McCleery, et al. 2014). Cities are viewed first in terms of their political value (where the voters are) rather than for the ecological values they may possess. The inherited historical view by general publics that urban environments are “biological deserts” seems reasonable as research has shown how sprawling urban development is responsible for high extinction rates (Luck 2007; McKinney & Schoch 2003: 298; McKinney 2008), extensive and persistent losses of native species (Hansen et al. 2005; Pickett et al. 1992) via large-scale transformation of landscapes (Ehrlich & Holdern 1971; Pejchar et al. 2007). However, urban ecology routinely necessitates re-assessing established ideas in biophysical ecology (e.g., linear responses of biodiversity to habitat destruction; Ramalho & Hobbs 2012; Collins et al. 2010; Grove et al. 2015) and advances in this field are transforming how we understand the ecological importance of our cities.

Such is the findings from the past decade of research on wild bees in cities. In the midst of a “pollination crisis” where insect pollinator populations are experiencing significant declines (Goulson et al. 2015; Jaffe et al. 2010; Pleasants & Oberhauser 2013), studies of native bee richness and abundance indicate diverse communities of wild bees persisting in cities in many parts of the world such as Berlin, Germany (Saure et al. 1998), Birmingham, Bristol, Cardiff, Dundee, Edinburg, Glasgow, Hull, Leeds, Leicester, London, Northampton, Reading, Sheffield, Southampton, and Swindon in the UK (Goulson et al. 2008; Baldock et al. 2015; Sirohi et al. 2015), Melbourne, Australia (Threfall et al. 2015), Guanacaste Province, Costa Rica (Frankie et al. 2013), Vancouver, Canada (Tommasi et al. 2004), and in the USA: Berkeley, CA (Frankie et al. 2005; 2016), Chicago, IL (Tonietto et al. 2011; Lowenstein et al. 2014), New York City, NY (Matteson et al. 2008; Matteson & Langellotto 2009), Phoenix, AZ (Cane et al. 2006), San

¹ We use “natural resource management” to broadly refer to nations’ governing bodies responsible for wildlife and land management such as a nation’s Ministry or Department of the Environment.

Francisco, CA (McFrederick & LeBuhn 2006), St. Louis, MO, and others. Bees found in these cities include both solitary and eusocial species, especially species that are cavity nesting and pollen generalists (Cariveau & Winfree 2015; Hernandez et al. 2009; Sirohi et al. 2015), even including specialized species indicative of high-quality habitat (e.g., pollen specialists and their cleptoparasites, Sheffield et al. 2013; Tonietto et al. 2011). In several cases, more diverse and abundant populations of native bees live in cities than in nearby rural landscapes (Baldock et al. 2015; Cane et al. 2006; Frankie et al., 2009; Matteson et al. 2008; Osborne et al. 2008; Verboven et al. 2014; Sirohi et al. 2015; for counter examples see Bates et al. 2011; Deguines et al. 2016; Geslin et al. 2013). For bumblebees in particular, urban areas can foster more species richness than rural or natural areas (Baldock et al. 2015; Gunnarsson & Federsel 2014; Winfree et al. 2007; McFrederick & LeBuhn 2006). Cities often contain greater bee species diversity than what would be expected from a more traditional viewpoint of urban areas.

Loss of habitat has been a long-term contributor to pollinator declines (Goulson et al. 2008; Harrison & Winfree 2015; Potts et al. 2010; Vanbergen 2013) while technological advances in agricultural efficiencies are increasingly homogenizing farmlands (Benton et al. 2003). Additional losses of natural areas to farming expansion and transition of traditional agricultural lands to those less hospitable to pollinators (e.g. monoculture commodity crops or indoor livestock operations) provides less floral forage over shorter periods of time (Ollerton et al. 2014; Scheper et al. 2014). Habitat loss and homogenization coupled with innovations of systemic pesticides and herbicides along with greater efficiency of chemical application have negatively affected wild pollinator populations in rural areas (Goulson et al. 2015; van der Sluijs et al. 2015; Simon-Delso et al. 2014; Straub et al. 2016; Whitehorn et al. 2012). While the protection and restoration of undeveloped lands are important for wild pollinator conservation and serve an obvious role in pollinator health, urban landscapes must not be overlooked. Surrounded by increasingly less hospitable rural and suburban landscapes², the city, with its variety of forage and nesting sites, becomes a refuge for insect pollinators.

² Relatively little is known about how urbanization affects ecological networks, but a few studies have emerged recently. Gotlieb et al. (2011; Central Jordan Rift Valley, Israel) and Geslin et al. (2013; Paris, France) on plant-

Advances in pollinator conservation in rural landscapes are proliferating across governance scales (President’s Task Force Strategy on Pollinator Health, 2015; Xerces Society; Pollinator Partnership, Intergovernmental Platform on Biodiversity & Ecosystem Services review, All-Ireland Pollinator Plan; Wales Pollinator Action Plan; others) but only a few governments are targeting urban landscapes and supporting these with funding (Natural Environment Research Council’s National Pollinator Strategy for England 2015; Welsh Action Plan for Pollinators; Living with Environmental Change Partnership). As urban ecology advances the science of ecology, the role of NRM agencies should similarly update their understanding of the role of cities in landscape-scale conservation priorities (see IPBES 2016). Engaging city planners and residents in enhancing insect pollinator habitat is a legitimate conservation practice in addition to its well-understood educational value. Implementing relevant programs requires collaborations, partnerships, and programming that reimagine the ecological value of urban lands: from “biological deserts” to valuable habitat for declining insect species.

This approach offers direct conservation benefits across a diversity of pollinator populations (cf. Kleijn et al. 2015) and the associated benefits across ecosystem services for humans (e.g. pollination of vegetables and fruit, cultural services associated with an interest in natural history, Peterson et al. 2010, others), plants (e.g. increased reproductive success), and animals (prey for higher trophic level species such as birds). Further, improving the wild pollinator populations in urban areas improve richness and abundance in nearby agricultural lands (Samnegård et al. 2011) via a spill-over effect (Goulson et al. 2010) though the relative importance of cities as sources or sinks for pollinators is largely unknown (Gill et al. 2016).

Intensifying conservation efforts for urban insect pollinators constitutes an opportunity for meaningful urban conservation—conservation that moves beyond traditional education and recreation

pollinator networks showed that the network became simplified with increased urbanization levels. A similar conclusion was reached by Rodewald et al. (2014) on plant-bird networks in Ohio, USA. In contrast, Sirohi et al. (in prep.) and Baldock et al. (2015) found that networks of flower visitors and their forage plants were no less complex in urban compared to rural settings, though there were significant differences in network structure and specialization of plants and insects, which was related to the larger number of plants in urban areas. The number of published studies is currently too limited, however, to make generalizations about how urbanization might affect plant-pollinator networks.

programming towards programming with cascading benefits throughout rural and urban landscapes. Matching conservation planning to the complexity of the city benefits NRM agencies via more direct connections to their constituency in population centers (Sanderson and Huron 2013). Conservation *for* the city finds an audience for agencies' other conservation efforts and likely, favor at the ballot box.

Pollinators put high-priority and high-impact urban conservation within reach. The relatively small spatial and temporal scales of insect pollinators in terms of functional ecology (habitat range, lifecycle, nesting behavior compared with larger mammals for example) offer opportunities for small actions to yield large benefits for pollinator health (cf. Frankie et al. 2014). The approach for improving the habitat value within urban areas is relatively simple and easily understood by urban residents. Several analyses and meta-analysis of urban insect pollinators found the consistent variable correlated with pollinator health is diverse forage—the presence of flowers (Bates et al. 2011; Cariveau and Winfree 2015; Hennig and Ghazoul 2012). These findings extend to forage species planted on urban vacant lands (Gardiner et al 2013) and rural lands (Scheper et al. 2013) with similar effects on specialist and generalist insect pollinators (Williams et al. 2010). Urban residential spaces play a role in pollinator abundance and diversity. Thus, individual decisions concerning yard management can have implications for conservation of threatened and endangered species (Goddard et al. 2010; Shwartz et al. 2013).

The city as refuge for insect pollinators opens many potential areas of research. Inventorying and monitoring is an essential practice to validate, improve, and communicate results of conservation efforts among partners and taxonomic experts. Understanding what works well and where is necessary for transferable practices across geographies which could aid decision makers across multiple scales. More research is needed to evaluate effectiveness of pollinator seed mixes (Garbuzov & Ratnieks 2014b), however, it is apparent that bees and other insect pollinators benefit from native and nonnative plants alike (da Silva Mouga 2015; Frankie et al. 2005; Hanley et al. 2014; Matteson & Langelotto 2011; Pardee & Philpott 2014), though for managerial purposes natives are preferred (Williams et al. 2011). Other underexplored areas include social dimensions of self-organizing neighbors transforming lawns

(and their affiliated cultural models) to attract bees and butterflies for conservation (Lerman & Milam 2016; van Heezik et al. 2012) and the effectiveness of various models as conservation strategies (Asah & Blahna 2013). Legal, political, and institutional questions regarding public land-uses and planting decision making, as well as institutional policies, organizational norms, and municipal ordinances affecting various actors' capacities of increasing pollinator habitat also require further investigation.

Cities offer several advantages for exploring conservation practices such as a lack of agriculture pesticides (Larson et al. 2013; Muratet and Fontaine 2015; even though "home" and horticultural pesticides may be widely used) and fewer large herbivores (e.g., deer) which allow some sensitive plants to be grown. Restoration work is fostered by relevant institutions, resources (e.g., museum collections), expert personnel (e.g., staff at botanical gardens), and volunteers who can install and maintain restoration plantings. Many of these urban resources are absent in rural areas. Cities also have concentrations of philanthropic donors, funding resources, and development specialists who can mobilize resources for conservation projects.

Coupling insect pollinator habitat enhancements with long-term species monitoring is one of the goals of the long-term wild bee monitoring being conducted in Chicago, IL, Detroit, MI, and St. Louis, MO (Tonietto et al. 2011). These projects are exploring social and cultural drivers of wild bee diversity and abundance in a variety of green spaces across these cities. Bee diversity in St. Louis, MO, seems to be responding to a combination of population density and income. For example, in low-income neighborhoods with low population density, bee diversity is higher than in denser and higher income ones (similar findings in Lowenstein et al. 2014; Lowenstein & Minor 2016). Not surprisingly, low-income, less-populated areas contain more vacant lots, and abandoned and crumbling infrastructure. Residential pesticide use is also reduced in low-income neighborhoods compared to higher income areas (Cook et al. 2012). More research is needed to understand the relationships between bee diversity and patterns of residential land-uses across shrinking and growing cities. Partnerships among city planners, citizen groups, conservation scientists, and policy makers targeting pollinator conservation can improve local

food security and community development. Improving global pollinator health across landscapes requires attending to populations of urban pollinators.

Research on urban insect pollinators is changing how we view the biological value and ecological importance of cities. Conservation must be repositioned within this newly unfolding image of the city. Rather than treating urban conservation as solely outreach and education aimed to improve political capital, NRM agencies can develop programming that improves natural capital thereby engaging urban citizens in improving the quality of life for threatened species living in cities. In 2050, it is estimated that 67% of the world's population will live in cities (UN 2014), much of these city landscapes have yet to be built (Grove et al. 2015). Attending to the conservation needs of insect pollinators among the suite of other green infrastructure, climate change adaptation, and environmental quality-of-life needs can inform current and future generations how to urbanize sustainably. To do so, requires an ecology *of* the city and its requisite conservation that fits the city: Conservation *for* the city.

Literature Cited

- Asah ST, Blahna DJ. 2013. Practical implications of understanding the influence of motivations on commitment to voluntary urban conservation stewardship. *Conservation Biology* **27**(4):866–875.
- Baldock KCR, et al. 2015. Where is the UK's pollinator biodiversity? The importance of urban areas for flower-visiting insects. *Proceedings of the Royal Society of London B: Biological Sciences* **282**(1803) DOI: 10.1098/rspb.2014.2849.
- Balooni K, Gangopadhyay K, Kumar BM. 2014. Governance for private green spaces in a growing Indian city. *Landscape and Urban Planning* **123**:21–29.
- Bates, AJ, Sadler, JP, Fairbrass, AJ, Falk, SJ, Hale, JD, Matthews, TJ. 2011. Changing bee and hoverfly pollinator assemblages along an urban-rural gradient. *PloS ONE* **6**(8): e23459.
- Benton TG, Vickery JA, Wilson JD. 2003. Farmland biodiversity: is habitat heterogeneity the key?. *Trends in Ecology & Evolution* **18**(4):182–188.
- Cane JH, Minckley RL, Kervin LJ, Roulston TAH, Williams NM. 2006. Complex responses within a desert bee guild (Hymenoptera: Apiformes) to urban habitat fragmentation. *Ecological Applications* **16**(2):632–644.
- Cariveau D, Winfree R. 2015. Causes of variation in wild bee responses to anthropogenic drivers. *Current Opinion in Insect Science* **10**:104–109.

- Carper AL, Adler LS, Warren PS, Irwin RE. 2014. Effects of suburbanization on forest bee communities. *Environmental Entomology* **43**(2):253–262.
- Collins SL, et al. 2010. An integrated conceptual framework for long-term social-ecological research. *Frontiers in Ecology and the Environment* **9**(6):351–357.
- Cook EM, Hall SJ, Larson KL. 2012. Residential landscapes as social-ecological systems: a synthesis of multi-scalar interactions between people and their home environment. *Urban Ecosystems* **15**(1):19–52.
- da Silva Mouga DMD, Feretti V, de Sena JC, Warkentin M, dos Santos AKG, Ribeiro CL. 2015. Ornamental bee plants as foraging resources for urban bees in Southern Brazil. *Agricultural Sciences* **6**:365–381.
- Deguines, N, Julliard, R, Flores, M, Fontaine, C. 2016. Functional homogenization of flower visitor communities with urbanization. *Ecology and evolution* **6**(7):1967–1976.
- Ehrlich PR, Holdren JP. 1971. Impact of population growth. *Science* **171**(3977):1212–1217.
- Frankie GW, Thorp RW, Schindler M, Hernandez J, Ertter B, Rizzardi M. 2005. Ecological patterns of bees and their host ornamental flowers in two northern California cities. *Journal of the Kansas Entomological Society* **78**(3):227–246.
- Frankie GW, Thorpe RW, Hernandez J, Rizzardi M., Ertter B, Pawelek JC, Witt SL, Schindler M, Coville R, Wojick VA. 2009. Native bees are a rich natural resource in urban California gardens. *California Agriculture* **63**(3):113–120.
- Frankie GW, Vinson SB, Rizzardi MA, Griswold TL, Coville RE, Grayum MH, Martinez LES, Foltz-Sweat J, Pawelek JC. 2013. Relationships of bees to host ornamental and weedy flowers in urban northwestern Guanacaste Province, Costa Rica. *Journal of Kansas Entomological Society* **86**(4):325–351.
- Frankie GW, Guerrero SL, Pawelek JC, Thorp R, Schindler M, Coville RE, Rizzaesi MA, Guinon M. 2016. Bees, flowers, and people in urban-agricultural-wildland interfaces. *Transactions of the 80th North American Wildlife and Natural Resources Management Conference*.
- Garbuzov M, Ratnieks FLW. 2014a. Quantifying variation among garden plants in attractiveness to bees and other flower-visiting insects. *Functional Ecology* **28**(2):364–374.
- Garbuzov M, Ratnieks FL. 2014b. Listmania: the strengths and weaknesses of lists of garden plants to help pollinators. *BioScience* DOI: 10.1093/biosci/biu150.
- Gardiner MM, Burkman CE, Prajzner SP. 2013. The value of urban vacant land to support arthropod biodiversity and ecosystem services. *Environmental Entomology* **42**(6):1123–1136.
- Geslin, B, Gauzens, B, Thébault, E, Dajoz, I. 2013. Plant pollinator networks along a gradient of urbanisation. *PLoS ONE* **8**(5):1–13.
- Gill RJ, Baldock KCR,... Potts, SG. 2016. Protecting an ecosystem service: approaches to understanding and mitigating threats to wild insect pollinators. *Advances in Ecological Research* **54**.

- Goddard M.A, Dougill AJ, Benton TG. 2010. Scaling up from gardens: biodiversity conservation in urban environments. *Trends in Ecology & Evolution* **25**(2):90–98.
- Goddard MA, Dougill AJ, Benton TG. 2013. Why garden for wildlife? Social and ecological drivers, motivations and barriers for biodiversity management in residential landscapes. *Ecological Economics* **86**:258–273.
- Gotlieb, A, Hollender, Y, Mandelik, Y. 2011. Gardening in the desert changes bee communities and pollination network characteristics. *Basic and Applied Ecology* **12**(4):310–320.
- Goulson D, Lye GC, Darvill B. 2008. Decline and conservation of bumble bees. *Annual Review of Entomology* **53**:191–208.
- Goulson D, Lepais O, O'Connor S, Osborne JL, Sanderson RA, Cussans J, Goffe L, Darvill B. 2010. Effects of land use at a landscape scale on bumblebee nest density and survival. *Journal of Applied Ecology* **47**(6):1207–1215.
- Goulson D, Nicholls E, Botías C, Rotheray EL. 2015. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science* **347**(6229) DOI: 10.1126/science.1255957.
- Grimm NB, Grove JM, Pickett STA, Redman CL. 2000. Integrated approaches to long-term studies of urban ecological systems. *BioScience* **50**(7):571–584.
- Grove JM, Locke DH, O'Neil-Dunne JPM. 2014. An ecology of prestige in New York City: Examining the relationships among population density, socio-economic status, group identity, and residential canopy cover. *Environmental Management* **54**(3):402–419.
- Grove JM, Cadenasso ML, Pickett STA, Machlis GE, Burch Jr. WR. 2015. *The Baltimore school of urban ecology: space, scale, and time for the study of cities*. Yale, New Haven.
- Gunnarsson B, Federsel LM. 2014. Bumblebees in the city: abundance, species richness and diversity in two urban habitats. *Journal of Insect Conservation* **18**(6):1185–1191.
- Hanley ME, Awbi AJ, Franco M. 2014. Going native? Flower use by bumblebees in English urban gardens. *Annals of Botany* **113**:799–806.
- Hansen AJ, Knight RL, Marzluff JM, Powell S, Brown K, Gude PH, Jones K. 2005. Effects of exurban development on biodiversity: patterns, mechanisms, and research needs. *Ecological Applications* **15**(6):1893–1905.
- Harrison T, Winfree R. 2015. Urban drivers of plant-pollinator interactions. *Functional Ecology* **29**(7):879–888.
- Hennig, EI, Ghazoul, J. 2012. Pollinating animals in the urban environment. *Urban Ecosystems* **15**(1):149–166.
- Hernandez JL, Frankie GW, Thorp RW. 2009. Ecology of urban bees: A review of current knowledge and directions for future study. *Cities and the Environment* **2**(1):3
- IPBES. 2016. Summary for policymakers of the assessment report of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services on pollinators, pollination and food

- production. S.G. Potts, V. L. Imperatriz-Fonseca, H. T. Ngo, J. C. Biesmeijer, T. D. Breeze, L. V. Dicks, L. A. Garibaldi, R. Hill, J. Settele, A. J. Vanbergen, M. A. Aizen, S. A. Cunningham, C. Eardley, B. M. Freitas, N. Gallai, P. G. Kevan, A. Kovács-Hostyánszki, P. K. Kwapong, J. Li, X. Li, D. J. Martins, G. Nates-Parra, J. S. Pettis, R. Rader, and B. F. Viana (eds.). pp. 1–28.
- Jaffe R, et al. 2010. Estimating the density of honeybee colonies across their natural range to fill the gap in pollinator decline censuses. *Conservation biology* **24**(2):583–593.
- Kleijn D, et al. 2015. Delivery of crop pollination services is an insufficient argument for wild pollinator conservation. *Nature Communications* **6** DOI: 10.1038/ncomms8414.
- Larson JL, Redmond CT, Potter DA. 2013. Assessing insecticide hazard to bumble bees foraging on flowering weeds in treated lawns. *PLoS ONE* **8**(6):1–7.
- Lerman SB, Milam J. 2016. Bee fauna and floral abundance within lawn-dominated suburban yards in Springfield, MA. *Annals of the Entomological Society of America* **109**(5):713–723.
- Lowenstein DM, Matteson KC, Xiao I, Silva AM, Minor ES. 2014. Humans, bees, and pollination services in the city: the case of Chicago, IL (USA). *Biodiversity and Conservation* **23**(11):2857–2874.
- Lowenstein, DM, Minor ES. 2016. Diversity in flowering plants and their characteristics: integrating humans as a driver of urban floral resources. *Urban Ecosystems* 1–14.
- Luck GW. 2007. A review of the relationships between human population density and biodiversity. *Biological Reviews* **82**(4):607–645.
- Matteson KC, Ascher JS, Langellotto GA. 2008. Bee richness and abundance in New York City urban gardens. *Annals of the Entomological Society of America* **101**(1):140–150.
- Matteson KC, Langellotto GA. 2009. Bumble bee abundance in New York City community gardens: implications for urban agriculture. *Cities and the Environment* **2**(1):5.
- Matteson, KC, Langellotto, GA. 2011. Small scale additions of native plants fail to increase beneficial insect richness in urban gardens. *Insect Conservation and Diversity* **4**(2):89–98.
- McFrederick QS, LeBuhn G. 2006. Are urban parks refuges for bumble bees *Bombus spp.* (Hymenoptera: Apidae)?. *Biological Conservation* **129**(3):372–382.
- McCleery RA, Moorman CE, Peterson MN. 2014. *Urban Wildlife Conservation: Theory and Practice*. Springer, New York.
- McKinney ML, Schoch RM, Yonavjak L. 2003. *Environmental science: Systems and solutions*. Jones & Bartlett Learning.
- McKinney ML. 2008. Effects of urbanization on species richness: a review of plants and animals. *Urban ecosystems* **11**(2):161–176.
- Muratet, A, Fontaine, B. 2015. Contrasting impacts of pesticides on butterflies and bumblebees in private gardens in France. *Biological Conservation* **182**:148–154.

- Ollerton J, Erenler H, Edwards M, Crockett R. 2014. Extinctions of aculeate pollinators in Britain and the role of large-scale agricultural changes. *Science* **346**(6215):1360–1362.
- Osborne JL, Martin AP, Shortall CR, Todd AD, Goulson D, Knight ME, Hale RJ, Sanderson RA. 2008. Quantifying and comparing bumblebee nest densities in gardens and countryside habitats. *Journal of Applied Ecology* **45**(3):784–792.
- Pardee GL, Philpott SM. 2014. Native plants are the bee’s knees: local and landscape predictors of bee richness and abundance in backyard gardens. *Urban Ecosystems* **17**(3):641–659.
- Pawelek J, Frankie GW, Thorp RW, Przybylski M. 2009. Modification of a community garden to attract native bee pollinators in urban San Luis Obispo, California. *Cities and the Environment* **2**(1):7.
- Pejchar L, Morgan PM, Caldwell MR, Palmer C, Daily GC. 2007. Evaluating the potential for conservation development: biophysical, economic, and institutional perspectives. *Conservation Biology* **21**(1): 69–78.
- Peterson, MJ, Hall, DM, Feldpausch, AM, Peterson, TR. 2010. Obscuring ecosystem function with application of the ecosystem services concept. *Conservation Biology* **24**(1):113–119.
- Pickett STA. 2012. Ecology of the city: a perspective from science. Pages 160–169 in urban design ecologies: AD reader. B. McGrath, ed. John Wiley and Sons Ltd, New York.
- Pickett STA, Parker VT, Fiedler PL. 1992. The new paradigm in ecology: implications for conservation biology above the species level. Pages 65–68 in Fielder PL, Jain SK, editors. *Conservation biology: the theory and practice of nature conservation preservation and management*. Springer, New York.
- Pleasants JM, Oberhauser KS. 2013. Milkweed loss in agricultural fields because of herbicide use: effect on the monarch butterfly population. *Insect Conservation and Diversity* **6**(2):135–144.
- Pollinator Health Task Force. 2015. National strategy to promote the health of honey bees and other pollinators, Washington, United States. Available from <https://www.whitehouse.gov/sites/default/files/microsites/ostp/Pollinator%20Health%20Strategy%202015.pdf> (accessed July 2015).
- Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. 2010. Global pollinator declines: trends, impacts and drivers. *Trends in Ecology & Evolution* **25**(6):345–353.
- Ramalho CE, Hobbs RJ. 2012. Time for a change: dynamic urban ecology. *Trends in Ecology & Evolution* **27**(3):179–188.
- Rodewald, AD, Rohr, RP, Fortuna, MA, Bascompte, J. 2014. Community-level demographic consequences of urbanization: an ecological network approach. *Journal of Animal Ecology* **83**(6):1409–1417.
- Samnegård, U, Persson, AS, Smith, HG. 2011. Gardens benefit bees and enhance pollination in intensively managed farmland. *Biological Conservation* **144**(11):2602–2606.
- Sanderson, EW, Huron A. 2011. Conservation in the city. *Conservation Biology* **25**(3):421–423.

- Saure C, Burger F, Dathe, HH. 1998. Die Bienenarten von Brandenburg und Berlin (Hym. Apidae). *Entomologische Nachrichten und Berichte* **42**(3):155–166.
- Sheffield CS, Pindar A, Packer L, Kevan PG. 2013. The potential of cleptoparasitic bees as indicator taxa for assessing bee communities. *Apidologie* **44**(5):501–510.
- Scheper J, Holzschuh A, Kuussaari M, Potts SG, Rundlöf M, Smith HG, Kleijn D. 2013. Environmental factors driving the effectiveness of European agri-environmental measures in mitigating pollinator loss—a meta-analysis. *Ecology letters*, **16**(7):912–920.
- Scheper J, Reemer M, van Kats R, Ozinga WA, van der Linden GTJ, Schaminée JHJ, Siepel H, Kleijn, D. 2014. Museum specimens reveal loss of pollen host plants as key factor driving wild bee decline in The Netherlands. *Proceedings of the National Academy of Sciences* **111**(49):17552–17557.
- Shwartz A, Muratet A, Simon L, Julliard R. 2013. Local and management variables outweigh landscape effects in enhancing the diversity of different taxa in a big metropolis. *Biological Conservation* **157**:285–292.
- Simon-Delso N, et al. 2014. Systemic insecticides (neonicotinoids and fipronil): trends, uses, mode of action and metabolites. *Environmental Science and Pollution Research* **22**(1):5–34.
- Sirohi MH, Jackson J, Edwards M, Ollerton J. 2015. Diversity and abundance of solitary and primitively eusocial bees in an urban centre a case study from Northampton (England). *Journal of Insect Conservation* **19**(3):487–500.
- Straub L, Villamar-Bouza L, Bruckner S, Chantawannakul P, Gauthier L, Khongphinitbunjong K, Retschnig G, Troxler A, Vidondo B, Neumann P, Williams GR. 2016. Neonicotinoid insecticides can serve as inadvertent insect contraceptives. *Proc. Royal Society B* **283**(1835):20160506.
- Threlfall CG, Walker K, Williams NSG, Hahs AK, Mata L, Stork N, Livesley SJ. 2015. The conservation value of urban green space habitats for Australian native bee communities. *Biological Conservation* **187**:240–248.
- Tommasi D, Miro A, Higo HA, Winston ML. 2004. Bee diversity and abundance in an urban setting. *The Canadian Entomologist* **136**(06):851–869.
- Tonietto R, Fant J, Ascher J, Ellis K, Larkin D. 2011. A comparison of bee communities of Chicago green roofs, parks and prairies. *Landscape and Urban Planning* **103**(1):102–108.
- Troy AR, Grove JM, O’Neil-Dunne JPM, Pickett STA, Cadenasso ML. 2007. Predicting opportunities for greening and patterns of vegetation on private urban lands. *Environmental Management* **40**(3):394–412.
- Tylianakis, JM, Laliberté, E, Nielsen, A, Bascompte, J. 2010. Conservation of species interaction networks. *Biological Conservation* **143**(10):2270–2279.
- United Nations. 2014. Our urbanizing world. Populations Facts: UN Department of Economic and Social Affairs **3**(August):1–5. http://www.un.org/en/development/desa/population/publications/pdf/popfacts/PopFacts_2014-3.pdf (accessed November 2015)

- 499 United States Fish & Wildlife Service. 2015. Urban wildlife conservation program, United States.
500 Available from: <http://www.fws.gov/urban/index.php> (accessed November 2015).
501
- 502 Van der Sluijs JP, et al. 2015. Conclusions of the Worldwide Integrated Assessment on the risks of
503 neonicotinoids and fipronil to biodiversity and ecosystem functioning. *Environmental Science*
504 *and Pollution Research* **22**(1):148–154.
505
- 506 van Heezik YM, Dickinson KJM, Freeman C. 2012. Closing the gap: communicating to change
507 gardening practices in support of native biodiversity in urban private gardens. *Ecology and*
508 *Society* **17**(1):34 DOI: 10.5751/ES-04712-170134.
509
- 510 Vanbergen AJ, et al. 2013. Threats to an ecosystem service: pressures on pollinators. *Frontiers in Ecology*
511 *and the Environment* **11**:251–259.
512
- 513 Verboven HAF, Uyttenbroeck R, Brys R, Hermy M. 2014. Different responses of bees and hoverflies to
514 land use in an urban-rural gradient show the importance of the nature of the rural land use.
515 *Landscape and Urban Planning* **126**:31–41.
516
- 517 Whitehorn PR, O'Connor S, Wackers FL, Goulson D. 2012. Neonicotinoid pesticide reduces bumble bee
518 colony growth and queen production. *Science* **336**(6079):351–352.
519
- 520 Williams NM, Crone EE, Roulston TAH, Minckley RL, Packer L, Potts SG. 2010. Ecological and life-
521 history traits predict bee species responses to environmental disturbances. *Biological*
522 *Conservation* **143**(10):2280–2291.
523
- 524 Williams NM, Cariveau D, Winfree R, Kremen C. 2011. Bees in disturbed habitats use, but do not prefer,
525 alien plants. *Basic and Applied Ecology* **12**(4):332–341.
- 526 Winfree R, Williams NM, Dushoff J, Kremen C. 2007. Native bees provide insurance against ongoing
527 honey bee losses. *Ecology Letters* **10**(11):1105–1113.